

Topological dimension of singular-hyperbolic attractors

C. A. Morales*

February 1, 2008

Abstract

An *attractor* is a transitive set of a flow to which all positive orbit close to it converges. An attractor is *singular-hyperbolic* if it has singularities (all hyperbolic) and is partially hyperbolic with volume expanding central direction [16]. The geometric Lorenz attractor [6] is an example of a singular-hyperbolic attractor with topological dimension ≥ 2 . We shall prove that *all* singular-hyperbolic attractors on compact 3-manifolds have topological dimension ≥ 2 . The proof uses the methods in [15].

1 Introduction

This paper is concerned with the topological dimension of attractors for flows on compact manifolds. By *attractor* we mean a transitive set of the flow to which all positive orbit close to it converges. The attractors under consideration will be *singular-hyperbolic* in the sense that they have singularities (all hyperbolic) and are partially hyperbolic with volume expanding central direction [16]. In particular, the singular-hyperbolic attractors are volume hyperbolic sets as defined in [1]. The geometric Lorenz attractor is an example of a singular-hyperbolic attractor with topological dimension ≥ 2 . We shall prove that *all* singular-hyperbolic attractors on compact 3-manifolds have topological dimension ≥ 2 . The proof uses the methods developed in [15]. Let us state our result in a precise way.

Hereafter X will be a C^1 vector field on a compact manifold M . The flow of X is denoted by X_t , $t \in \mathbb{R}$. Given $p \in M$ we define $\omega(p) = \omega_X(p)$, the ω -limit set of p , as the accumulation point set of the positive orbit of p . The α -limit set of p

*2000 MSC: Primary 37D30, Secondary 37C45. *Key words and phrases:* Attractor, Partially Hyperbolic, Topological Dimension. Partially supported by CNPq, FAPERJ and PRONEX-Dyn. Sys./Brazil.

is the set $\alpha(p) = \alpha_X(p) = \omega_{-X}(p)$. A compact invariant set Λ of X is *transitive* or *attracting* depending on whether $\Lambda = \omega(p)$ or $\cap_{t>0} X_t(U)$ for some $p \in \Lambda$ or some compact neighborhood U of Λ respectively. An *attractor* is a transitive attracting sets. A closed orbit of X is either periodic or singular. A singularity of X is *hyperbolic* if none of its eigenvalues have zero real part.

A compact invariant set Λ of X is *partially hyperbolic* [8] if there are an invariant splitting $T\Lambda = E^s \oplus E^c$ and positive constants K, λ such that:

1. E^s is contracting, namely

$$\|DX_t/E_x^s\| \leq Ke^{-\lambda t}, \quad \forall x \in \Lambda, \forall t > 0.$$

2. E^s dominates E^c , namely

$$\|DX_t/E_x^s\| \cdot \|DX_{-t}/E_{X_t(x)}^c\| \leq Ke^{-\lambda t}, \quad \forall x \in \Lambda, \forall t > 0.$$

The central direction E^c of Λ is said to be *volume expanding* if the additional condition

$$|\text{Det}(DX_t/E_x^c)| \leq Ke^{-\lambda t}$$

holds $\forall x \in \Lambda, \forall t > 0$ where $\text{Det}(\cdot)$ means the jacobian. The above splitting $E^s \oplus E^c$ will be referred to as a (K, λ) -splitting in the Appendix.

Definition 1.1. ([16]) *An attractor is singular-hyperbolic if it has singularities (all hyperbolic) and is partially hyperbolic with volume expanding central direction.*

Definition 1.2. ([9]) *The topological dimension of a space E is either -1 (if $E = \emptyset$) or the last integer k for which every point has arbitrarily small neighborhoods whose boundaries have dimension less than k .*

The relation between dynamics and topological dimension was considered for hyperbolic systems [7, 18, 2, 3]; for expansive systems [10, 13]; and for singular-hyperbolic systems [14]. The result below generalizes to singular-hyperbolic attractors a well known property of both hyperbolic strange attractors and geometric Lorenz attractors.

Theorem A . *Singular-hyperbolic attractors on compact 3-manifolds have topological dimension ≥ 2 .*

The idea of the proof is the following. Let Λ be a singular-hyperbolic attractor of a flow X on a compact 3-manifold M . It follows from [16] that all the singularities $\sigma \in \Lambda$ are *Lorenz-like*, namely the eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of σ are real and satisfy $\lambda_2 < \lambda_3 < 0 < -\lambda_3 < \lambda_1$. The flow nearby σ can be described using the Grobman-Hartman Theorem [5]. In particular, a Lorenz-like singularity exhibits

two *singular cross-sections* S^t, S^b and two *singular curves* l^t, l^b ([15]). A *singular cross section* of Λ is by definition a disjoint collection of singular cross sections S^t, S^b (as σ runs over all the singularities of Λ) whose horizontal boundaries does not intersect Λ . The *singular curve* of S is the union l of the respective singular curves l^t, l^b . A *singular partition* of Λ will be a compact neighborhood O of $\Lambda \cap l$ in S , for some singular cross section S of Λ , such that $\Lambda \cap l$ does not intersect the boundary of O and every regular orbit of Λ intersect O . The *size* of the singular partition O is the minimal $\epsilon > 0$ such that there is an invariant cone field in O (for the return map $\Pi : \text{Dom}(\Pi) \subset O \rightarrow O$) on which the derivative of Π has expansion rate bigger than ϵ^{-1} . In Proposition 2.7 we prove that one-dimensional singular-hyperbolic attractors on compact 3-manifolds have singular partition with arbitrarily small size. The proof of this proposition uses the Lemmas 7.5 and 7.6 in [15]. These lemmas will be proved in the Appendix for the sake of completeness. In Theorem 2.9 we shall prove that singular-hyperbolic attractors Λ on compact 3-manifolds cannot have singular partitions with arbitrarily small size. Theorem A will follow from Proposition 2.7 and Theorem 2.9.

2 Proof

We start with some definitions. Hereafter Λ is a singular-hyperbolic attractor of a C^1 flow X on a compact 3-manifold M . Since Lorenz-like singularities σ are hyperbolic they are equipped with three invariant manifolds $W_X^s(\sigma), W_X^u(\sigma), W_X^{ss}(\sigma)$ each one tangent at σ to the eigenspace corresponding to $\{\lambda_2, \lambda_3\}, \{\lambda_1\}, \{\lambda_2\}$ respectively. It follows from [16] that every singularity σ of X in Λ is Lorenz-like and satisfies $\Lambda \cap W_X^{ss}(\sigma) = \{\sigma\}$. The classical Grobman-Hartman Theorem [5] gives the description of the flow nearby σ . This is done at Figure ???. Note that $W_X^{ss}(\sigma)$ separates $W_X^s(\sigma)$ in two connected components denoted the top and the bottom respectively. In one of these components, say the top one, we consider a cross-section $S^t = S_\sigma^t$ together with a curve $l^t = l_\sigma^t$ as in Figure ??. Similarly we consider a cross-section $S^b = S_\sigma^b$ and a curve $l^b = l_\sigma^b$ located in the bottom component of $W_X^s(\sigma)$. See Figure ??. Both S^* (for $*$ = 1, 2) are homeomorphic to $[0, 1] \times [0, 1]$. S^* can be chosen in a way that l^* is contained in $W_X^s(\sigma) \setminus W_X^{ss}(\sigma)$. The positive flow lines of X starting at $S^t \cup S^b \setminus (l^t \cup l^b)$ exit a small neighborhood of σ passing through the cusp region as indicated in Figure ??. The positive orbits starting at $l^t \cup l^b$ goes directly to σ . We note that the boundary of S^* is formed by four curves, two of them transverse to l^* and two of them parallel to l^* . The union of the curves in the boundary of S^* which are parallel (resp. transverse) to l^* is denoted by $\partial^v S^*$ (resp. $\partial^h S^*$). The interior (as a submanifold) of S^* is denoted by $\text{Int}(S^*)$.

Remark 2.1. *An immediate consequence of $\Lambda \cap W_X^{ss}(\sigma) = \{\sigma\}$ is the following. Let σ be a singularity of X in Λ . Then there are cross-sections S^t, S^b as above arbitrarily close to σ such that $\Lambda \cap \partial^h S^* = \emptyset$ ($* = t, b$). Since the two boundary points of l^* are in $\partial^h S^*$ we have that $\Lambda \cap l^* \subset \text{Int}(S^*)$.*

Definition 2.2. *We shall call the cross sections S^t, S^b as singular cross sections associated to σ . The curves l^t, l^b are called singular curves of S^t, S^b respectively. A singular cross section of Λ is a finite disjoint collection $\{S_\sigma^t, S_\sigma^b : \sigma \text{ is a singularity of } X \text{ in } \Lambda\}$ satisfying $\Lambda \cap \partial^h S = \emptyset$. The singular curve of S is the associated collection of singular curves $l = \{l_\sigma^t, l_\sigma^b : \sigma \text{ is a singularity of } X \text{ in } \Lambda\}$.*

Hereafter we denote by $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$ the singular-hyperbolic splitting of Λ . The contracting direction E^s is one-dimensional and contracting. So, E_Λ^s can be extended to an invariant contracting splitting $E_{U(\Lambda)}^s$ on a neighborhood $U(\Lambda)$ of Λ . The standard Invariant Manifold Theory [8] implies that $E_{U(\Lambda)}^s$ is tangent to a continuous foliation $\mathcal{F}$ on $U(\Lambda)$. If S is a singular cross-section contained in $U(\Lambda)$, we denote by $\mathcal{F}^S$ the foliation of S obtained projecting $\mathcal{F}$ into S along X . The space of leaves of $\mathcal{F}^S$ will be denoted by I^S . We extend E_Λ^c continuously to a subbundle $E_{U(\Lambda)}^c$ of $T_{U(\Lambda)}M$. In what follows we fix such a neighborhood $U(\Lambda)$ of Λ .

Remark 2.3. *It is possible to choose S arbitrarily close to the singularities of Λ in a way that l is a finite union of leaves of $\mathcal{F}^S$ and I^S is a finite disjoint union of compact intervals.*

The following lemma is a direct consequence of standard argument involving topological dimension. We prove it here for the sake of completeness.

Lemma 2.4. *Let S a singular cross-section and l be its associated singular curve. If Λ is one-dimensional, then there is a compact neighborhood O of $\Lambda \cap l$ in S whose boundary ∂O satisfies $\Lambda \cap \partial O = \emptyset$.*

Proof. Note that $\Lambda \cap \partial^h S = \emptyset$ since S is a singular cross-section. As noted in Remark 2.1 one has $\Lambda \cap l \subset \text{Int}(S)$. Fix $x \in \Lambda \cap l$. Then $x \in \text{Int}(S)$. Because Λ is one dimensional we have that $\Lambda \cap S$ is zero dimensional [9]. Then, by the definition of the topological dimension, one can find an open set S_x of $\Lambda \cap S$ containing x such that $\partial S_x = \emptyset$. Note that the topology in $\Lambda \cap S$ is the one induced by S . It follows that $S_x = (\Lambda \cap S) \cap O_x$ for some open set O_x of M . Since S is transversal to X we can choose O_x such that $\partial S_x = (\Lambda \cap S) \cap \partial O_x$ (for this we can use the Tubular Flow-Box Theorem [5]). It follows that

$$(\Lambda \cap S) \cap \partial O_x = \emptyset.$$

On the other hand, $\Lambda \cap l$ is compact in S and $\{S \cap O_x : x \in \Lambda \cap l\}$ is an open covering of $\Lambda \cap l$. It follows that there is a finite subcollection of $\{S \cap O_x : x \in \Lambda \cap l\}$

covering $\Lambda \cap l$. Denote by O the union of the closures (in S) of the elements of such a subcollection. It follows that O is a compact neighborhood of $\Lambda \cap l$ in S . Since O is a finite union of $S \cap O_x$'s satisfying $(\Lambda \cap S) \cap \partial O_x = \emptyset$ we have that $\Lambda \cap \partial O = \emptyset$. This proves the lemma. $\square$

Hereafter O is a set contained in a singular cross section S . Clearly O defines a return map

$$\Pi : \text{Dom}(\Pi) \subset O \rightarrow O$$

given by

$$\Pi(x) = X_{t(x)}(x),$$

where $\text{Dom}(\cdot)$ denotes the domain and $t(\cdot)$ denotes the return time.

Remark 2.5. *Note that Π may be discontinuous in $\Pi^{-1}(\partial O)$. However if $x \in \Pi^{-1}(\text{Int}(O))$ then Π is C^1 in an open neighborhood of x contained in $\text{Int}(O)$. This is an immediate consequence of the Tubular Flow-Box Theorem.*

We denote by TO the tangent space of O relative S . If $x \in M$ we denote by $\angle(v_x, w_x)$ the tangent of the angle between $v_x, w_x \in T_x M$. If L_x is a linear subspace of $T_x M$, we define

$$\angle(v_x, L_x) = \inf_{w_x \in L_x} \angle(v_x, w_x).$$

Given $\alpha > 0$ we define the *cone*

$$C_\alpha(L_x) = \{v_x \in T_x M : \angle(v_x, L_x) \leq \alpha\}.$$

If $L : x \in \text{Dom}(L) \rightarrow L_x$ is a map and $\alpha > 0$ we define the *cone field*

$$\mathcal{C}_\alpha(L) = \{C_\alpha(L_x) : x \in \text{Dom}(L)\}.$$

The case $L = E^c$ will be interesting. The definition below is a minor modification of the corresponding definition in [15]. If $x \in M$ we denote $X_{\mathbb{R}}(x)$ the full orbit of x .

Definition 2.6. *A singular partition of Λ is a set O satisfying the following properties:*

1. *There is a singular cross-section S such that $O \subset \text{Int}(S)$ is a compact neighborhood of $\Lambda \cap l$.*
2. $\Lambda \cap \partial O = \emptyset$.
3. $\text{Sing}_X(\Lambda) = \{q \in \Lambda : X_{\mathbb{R}}(q) \cap O = \emptyset\}$.

The size of O is the minimal number $\epsilon > 0$ for which there is $\alpha > 0$ such that the cone field $\mathcal{C}_\alpha(E^c)$ satisfies :

4. If $x \in \text{Dom}(\Pi)$, then

$$D\Pi(x)(C_\alpha(E_x^c) \cap TO_x) \subset \text{int}(C_{\frac{\alpha}{2}}(E_{\Pi(x)}^c) \cap TO_{\Pi(x)}).$$

5. If $x \in \text{Dom}(\Pi)$ and $v_x \in C_\alpha(E_x^c) \cap TO_x$, then

$$\|D\Pi(x)(v_x)\| \geq \epsilon^{-1}\|v_x\|.$$

6. $\inf\{\angle(v_x, E_x^s) : x \in O, v_x \in C_\alpha(E_x^c) \cap TO_x\} > 0$.

The following proposition studies the existence of singular partition with arbitrarily small size for certain one-dimensional singular-hyperbolic sets. Its proof uses the methods developed in [15]. We let $\text{Sing}_X(\Lambda) = \{\sigma_1, \dots, \sigma_k\}$ be the set of singularities of X in Λ .

Proposition 2.7. *One-dimensional singular-hyperbolic attractors on compact 3-manifolds have singular partitions with arbitrarily small size.*

Proof. Let Λ be a singular-hyperbolic attractor of a C^1 flow X on a compact 3-manifold M . We shall assume that Λ has topological dimension 1. We shall prove that Λ has singular partition with arbitrarily small size $\epsilon > 0$. For this we proceed as follows. Since Λ has topological dimension 1 we have that Λ cannot contain hyperbolic sets (the unstable manifold of a hyperbolic set in Λ would be two-dimensional and contained in Λ). It follows that $\omega(x)$ cannot be hyperbolic for all $x \in \Lambda$. By [17] if $\mathcal{L} = \omega(x), \alpha(x)$ then

$$\mathcal{L} \cap \text{Sing}_X(\Lambda) \neq \emptyset, \quad \forall x \in \Lambda. \quad (1)$$

Choose $\alpha > 0$ such that

$$\inf\{\angle(v_x, E_x^s) : x \in U(\Lambda), v_x \in C_\alpha(E_x^c)\} > 0.$$

By [15, Lemma 7.5] (see Lemma 3.1) we can find a neighborhood $U_\alpha \subset U(\Lambda)$ of Λ and positive constants $T_\alpha, K_\alpha, \lambda_\alpha$ such that the following properties hold:

(P1). If $x \in U_\alpha$ and $t \geq T_\alpha$, then

$$DX_t(x)(C_\alpha(E_x^c)) \subset C_{\alpha/2}(E_{X_t(x)}^c).$$

(P2). If $x \in U_\alpha$ is regular, $X(x) \in C_\alpha(E_x^c)$, $t \geq T_\alpha$ and $v_x \in C_\alpha(E_x^c)$ is orthogonal to $X(x)$, then

$$\|P_x^t(v_x)\| \cdot \|X(X_t(x))\| \geq K_\alpha e^{\lambda_\alpha t} \cdot \|v_x\| \cdot \|X(x)\|,$$

where P_x^t denotes the *Poincaré flow associated to X* (see [4, 15] or the Appendix).

Once we fix α and U_α we apply [15, Lemma 7.6] (see Lemma 3.2) to find, for every $i \in \{1, \dots, k\}$, a pair of singular cross-sections $S_i^{*,0}$ associated to σ_i ($* = t, b$) such that

$$X(x) \in C_\alpha(x), \quad \forall x \in S_i^{*,0}.$$

Define

$$S^0 = \cup_{i=1}^k (S_i^{t,0} \cup S_i^{b,0}).$$

It is clear that S^0 is a singular cross-section. We denote by l^0 the singular curve of S^0 . Since S^0 is transversal to X one can find a constant $D > 0$ (depending on S^0) such that

$$\frac{\|X(x)\|}{\|X(y)\|} > D,$$

for all $x, y \in S^0$. We choose $T_\epsilon > T_\alpha$ large enough so that

$$K_\alpha e^{\lambda_\alpha t} \cdot D > \epsilon^{-1} \quad (2)$$

for all $t \geq T_\epsilon$.

For every $\delta > 0$ we consider a singular-cross section $S^\delta \subset S^0$ ($i = 1, \dots, k$ and $* = t, b$) formed by small bands $S_i^{*,\delta}$ of diameter 2δ around the singular curve l_i^* of $S_i^{*,0}$. Note that the singular curve of S^δ is l^0 (the one of S^0) for all δ . Since Λ is one-dimensional Lemma 2.4 implies that $\forall x \in \Lambda \cap l^0$ there is a compact neighborhood $O = O^\delta \subset S^\delta$ of $\Lambda \cap l^0$ such that $\Lambda \cap \partial O = \emptyset$. Note that O is a singular partition of Λ . In fact, (1) and (2) of Definition 2.6 are obvious. And (3) of Definition 2.6 follows from Eq. (1) since O is a compact neighborhood of $\Lambda \cap l^0$.

Let us prove that if $\delta > 0$ small enough, then O has size ϵ . For this we need to prove that for δ small the cone field $C_\alpha(E^c)$ satisfies the properties (4)-(6) of Definition 2.6. Let $\Pi : \text{Dom}(O) \subset O \rightarrow O$ be the return map induced by X in O . By definition $\Pi(x) = X_{t(x)}(x)$ where $t(x)$ is the return time of $x \in \text{Dom}(O) \subset O$ into O . To calculate $D\Pi(x)$ we can assume without loss of generality that S^0 is orthogonal to X . It follows that

$$D\Pi(x) = P_x^{t(x)}$$

for all $x \in \text{Dom}(\Pi)$. Shrinking δ one has $t(x) > T_\epsilon$ for all $x \in \text{Dom}(\Pi) \subset O$. This allows us to apply the properties (P1)-(P2) above. In fact, since $D\Pi(x) = P_x^{t(x)}$ one has

$$D\Pi(x)/C_\alpha(E_x^c) \cap TO_x = P_x^{t(x)}/C_\alpha(E_x^c) \cap TO_x.$$

Then, Definition 2.6-(4) follows from (P1). (P2) and Eq. (2) imply

$$\|D\Pi(x)(v_x)\| = \|P_x^{t(x)}(v_x)\| =$$

$$\begin{aligned}
&= \|P_x^{t(x)}(v_x)\| \cdot \|X(X_{t(x)}(x))\| \cdot \|X(X_{t(x)}(x))\|^{-1} \geq \\
&\geq K_\alpha e^{\lambda_\alpha t(x)} \cdot \frac{\|X(x)\|}{\|X(X_{t(x)}(x))\|} \cdot \|v_x\| \geq K_\alpha e^{\lambda_\alpha t(x)} \cdot D \cdot \|v_x\| \geq \epsilon^{-1} \|v_x\|,
\end{aligned}$$

$\forall x \in \text{Dom}(\Pi)$, $\forall v_x \in C_\alpha(E_x^c) \cap TO_x$ because $X(x) \in C_\alpha(E_x^c) \forall x \in S^0$ and $t(x) > T_\epsilon$ ($\forall x \in \text{Dom}(\Pi)$). This proves Definition 2.6-(5). Definition 2.6-(6) is a direct consequence of the choice of α . The result follows. $\square$

The following lemma will be used to prove Theorem 2.9. Recall that if S is a singular cross-section then S is endowed with a singular curve l . If $O \subset S$ then $\Pi : \text{Dom}(\Pi) \subset O \rightarrow O$ denotes the return map associated to O .

Lemma 2.8. *Let Λ be a singular-hyperbolic attractor of a flow X on a compact 3-manifold. Let O be a singular partition of Λ . Then, there is an open neighborhood $O' \subset O$ of $\Lambda \cap O$ such that:*

1. $O' \setminus l \subset \text{Dom}(\Pi)$.
2. Π is C^1 in $O' \setminus l$.
3. $\Pi(O' \setminus l) \subset O'$.

Proof. Because Λ is an attractor we have that the unstable manifold of any of its singularities is contained in Λ . In particular, every connected component of $W_X^u(\sigma_i) \setminus \{\sigma_i\}$ is contained in $\Lambda \forall i$. It follows from Definition 2.6-(3) that all such components intersect O . By Definition 2.6-(2) such intersections can occur only in $\text{Int}(O)$. This implies that there are small open bands, centered at the singular curves in l , whose union $V(l)$ satisfies $V(l) \setminus l \subset \Pi^{-1}(\text{Int}(O))$. As noted in Remark 2.5 we have that Π is C^1 in $V(l) \setminus l$. Again by Definition 2.6-(2)-(3) one has $(\Lambda \cap O) \setminus V(l) \subset \Pi^{-1}(\text{Int}(O))$. So, by Remark 2.5, since $(\Lambda \cap O) \setminus V(l)$ is compact, there is an open neighborhood V of $(\Lambda \cap O) \setminus V(l)$ contained in $\text{Dom}(\Pi)$ such that Π is C^1 in V . Observe that $V \cup V(l)$ is an open neighborhood of $\Lambda \cap O$ such that Π is C^1 in $(V \cup V(l)) \setminus l$. On the other hand, Λ is an attractor by assumption. Then, there is a neighborhood U^* such that $X_t(U^*) \subset U^*, \forall t > 0$. Clearly one can choose U^* to be *arbitrarily close to* Λ . In particular, $O' := O \cap U^*$ is contained in $V \cup V(l)$. It follows that $O' \setminus l \subset \text{Dom}(\Pi)$ because $V \cup (V(l) \setminus l) \subset \text{Dom}(\Pi)$. Because $X_t(U^*) \subset U^*$ for all $t > 0$ and the return time for the points in $\text{Dom}(\Pi)$ is positive we conclude that $\Pi(O' \setminus l) \subset O'$. As Π is C^1 in $(V \cup V(l)) \setminus l$ and $O' \subset V \cup V(l)$ we conclude that Π is C^1 in $O' \setminus l$. This proves the result. $\square$

Theorem 2.9. *Singular-hyperbolic attractors on compact 3-manifolds cannot have singular partitions with arbitrarily small size.*

Proof. Let Λ be a singular-hyperbolic attractor of a C^1 flow X on a compact 3-manifold M . By contradiction we assume that Λ has a singular partition O with arbitrarily small size $\epsilon > 0$. We fix $\epsilon \in (0, 1/2)$. We let O' be the open neighborhood obtained in Lemma 2.8 for O . Hereafter we say that a C^1 connected curve c in O is a C^u -curve if its tangent vector belongs to the cone field $C_\alpha(E^c)$ at Definition 2.6-(4). Definition 2.6-(4) implies that Π carries C^u curves in $\Pi^{-1}(Int(O))$ into C^u curves in O (see also Remark 2.5). Definition 2.6-(6) implies that a C^u curve in O intersects l in at most one point x_c . In that case x_c divides c in two connected components the largest one being denoted by c^+ . Clearly if $L(\cdot)$ denotes the lenght, then

$$L(c^+) \geq (1/2)L(c).$$

Now, fix a C^u curve $c_1 \subset O' \setminus l$. Define $R = (2\epsilon)^{-1}$. The choice of ϵ implies $\epsilon^{-1} > R > 1$. Lemma 2.8-(1) implies $c_1 \subset \text{Dom}(\Pi)$. Lemma 2.8-(2) implies that $c_2 = \Pi(c_1)$ is a C^u curve contained in O' . Definition 2.6-(5) implies $L(c_2) \geq \epsilon^{-1}L(c_1) \geq R \cdot L(c_1)$. Suppose we have constructed a sequence $c_1, c_2, \dots, c_i$ of C^u curves of O contained in O' satisfying $L(c_j) \geq R \cdot L(c_{j-1})$ for all $2 \leq j \leq i$. If $c_i \cap l = \emptyset$ we define $c_{i+1} = \Pi(c_i)$ and keep going. If $c_i \cap l \neq \emptyset$ we define $c_{i+1} = \text{Closure}(\Pi(c_i^+))$. In any case c_{i+1} is a C^u curve of O contained in O' . In the first case we have $L(c_{i+1}) \geq \epsilon^{-1}L(c_i) \geq R \cdot L(c_i)$. In the second case we have

$$L(c_{i+1}) = L(\Pi(c_i^+)) \geq \epsilon^{-1}L(c_i^+) \geq (\epsilon^{-1}/2) \cdot L(c_i) = R \cdot L(c_i).$$

In this way we can construct an infinite sequence $c_1, \dots, c_i, c_{i+1}, \dots$ of C^u curves of O in O' all of which satisfying $L(c_{i+1}) \geq R \cdot L(c_i)$. It follows that

$$L(c_i) \geq R^i \cdot L(c_1),$$

for all i . Since $l(c_1) > 0$ and $R = (2\epsilon)^{-1} > 1$ we conclude that

$$\lim_{i \rightarrow \infty} L(c_i) = \infty.$$

On the other hand, let S be the singular cross-section containing O given by Definition 2.6-(1). Let $\mathcal{F}^S$ be the projection of the stable manifold in $U(\Lambda)$ over S . As noted in Remark 2.3 the leave space I^S of $\mathcal{F}^S$ is a finite union of compact intervals. In particular I^S has finite diameter. Since $O' \subset O \subset Int(S)$ we have that all the curves c_i are contained in S . Since c_i is a C^u curve we have by Definition 2.6-(6) that c_i have positive angle with the leaves of $\mathcal{F}^S$ (note that these leaves are tangent to E^s). So, we can project c_i to obtain an infinite sequence of intervals in I^S . The lenght of these intervals goes to ∞ (as $i \rightarrow \infty$) since $L(c_i) \rightarrow \infty$ (as $i \rightarrow \infty$). This is a contradiction since I^S has finite diameter. This contradiction proves the result. $\square$

Proof of Theorem A : Let Λ be a singular-hyperbolic attractor on a compact 3-manifold. If Λ has topological dimension < 2 then Λ would be one-dimensional because it has regular orbits [9]. It would follow from Proposition 2.7 that Λ has singular partitions with arbitrarily small size contradicting Theorem 2.9. The proof follows. $\square$

3 Appendix

In this section we state (and prove) two technical lemmas which were used in the proof of Theorem A. These lemmas were proved in [15] and here we reproduce these proofs for the sake of completeness. Let us state some definitions and notations.

First we define the Linear Poincaré Flow [4]. Let X be a flow on a compact 3-manifold M . The Riemmanian Metric of M is denoted by $\langle \cdot, \cdot \rangle$. If x is a regular point of X (i.e. $X(x) \neq 0$), we denote by $N_x = \{v_x \in T_x M : \langle v_x, X(x) \rangle = 0\}$ the orthogonal complement of $X(x)$ in $T_x M$. Denote $O_x : T_x M \rightarrow N_x$ the orthogonal projection onto N_x . For every $t \in \mathbb{R}$ we define $P_x^t : N_x \rightarrow N_{X_t(x)}$ by

$$P_x^t = O_{X_t(x)} \circ DX_t(x).$$

It follows that $P = \{P_x^t : t \in \mathbb{R}, X(x) \neq 0\}$ satisfies the cocycle relation

$$P_x^{s+t} = P_{X_t(x)}^s \circ P_x^s,$$

for every $t, s \in \mathbb{R}$. The parametrized family P is called the *Linear Poincaré Flow* of X .

We denote by $\text{vol}(v_x, w_x)$ the area of the parallelogram in $T_x M$ generated by $v_x, w_x \in T_x M$. As M is a compact manifold, there is a constant $V \geq 1$ such that $V^{-1} \leq \text{vol}(v_x, w_x) \leq V$, $\forall x \in M, \forall v_x, w_x \in T_x M$ satisfying $\|v_x\| = \|w_x\| = 1$ and $\langle v_x, w_x \rangle = 0$. For simplicity we shall assume that $V = 1$. In other words,

$$\text{vol}(v_x, w_x) = \|v_x\| \cdot \|w_x\|,$$

$\forall x \in M, \forall v_x, w_x \in T_x M$ with $\langle v_x, w_x \rangle = 0$.

In addition,

$$\text{vol}(v_x, X(x)) = \|O_x v_x\| \cdot \|X(x)\|,$$

$\forall x \in M$ regular, $\forall v_x \in T_x M$. In particular,

$$\text{vol}(DX_t(x)v_x, X(X_t(x))) = \|P_x^t(v_x)\| \cdot \|X(X_t(x))\|, \quad (3)$$

$\forall x \in M$ regular, $\forall t \in \mathbb{R}, \forall v_x \in N_x$.

Recall that if Λ is a singular hyperbolic set of X with (K, λ) -splitting $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$, then

$$| \text{Det}(DX_t/E_x^c) | \geq K e^{\lambda t},$$

$\forall x \in \Lambda, \forall t \geq 0$, where $\text{Det}(\cdot)$ denotes the jacobian. So, if Λ is a singular hyperbolic set as above one has

$$\text{vol}(DX_t(x)v_x^c, DX_t(x)w_x^c) \geq K e^{\lambda t} \text{vol}(v_x^c, w_x^c),$$

$\forall x \in \Lambda, \forall t \geq 0, \forall v_x^c, w_x^c \in E_x^c$.

Remember that $U(\Lambda)$ denotes a neighborhood of Λ where the splitting $E_\Lambda^s \oplus E_\Lambda^c$ extends to $E_{U(\Lambda)}^s \oplus E_{U(\Lambda)}^c$.

Lemma 3.1. *Let Λ be a singular-hyperbolic attractor of a C^1 flow X on a compact 3-manifold M . Then for every $\alpha \in (0, 1]$ there are a neighborhood $U_\alpha \subset U(\Lambda)$ of Λ and constants $T_\alpha, K_\alpha, \lambda_\alpha > 0$ such that :*

1. *If $x \in U_\alpha$ and $t \geq T_\alpha$, then*

$$DX_t(x)(C_\alpha(E_x^c)) \subset C_{\frac{\alpha}{2}}(E_{X_t(x)}^c).$$

2. *If $x \in U_\alpha$ is regular, $X(x) \in C_\alpha(E_x^c)$, $t \geq T_\alpha$ and $v_x \in C_\alpha(E_x) \cap N_x$, then*

$$\|P_x^t(v_x)\| \cdot \|X(X_t(x))\| \geq K_\alpha e^{\lambda_\alpha t} \cdot \|v_x\| \cdot \|X(x)\|.$$

Proof. Let Λ and $\alpha \in (0, 1]$ be as in the statement. As mentioned above $T_{U(\Lambda)} M = E_{U(\Lambda)}^s \oplus E_{U(\Lambda)}^c$ denotes the extension of the (K, λ) -splitting $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$ of Λ to a neighborhood $U(\Lambda)$ of Λ . Let π^s the projection of T_Λ on E_Λ^s , and π^c be the projection of T_Λ on E_Λ^c . Denote $v_x = v_x^s + v_x^c \in E_x^s \oplus E_x^c = T_x M \forall x \in U(\Lambda)$, $\forall v_x \in T_x M$. In other words, $v_x^s = \pi^s(v_x)$ and $v_x^c = \pi^c(v_x)$

As E_Λ^s (K, λ) -dominates E_Λ^c we have that

$$\|DX_t(x)/E_x^s\| \leq K^{-1} e^{-\lambda t} m(DX_t(x)/E_x^c), \quad (4)$$

$\forall x \in \Lambda, \forall t \geq 0$.

Fix $R > 4$ such that

$$\frac{K}{R} < 1. \quad (5)$$

Choosing $T^1 = T_\alpha^1 > 0$ large enough one has

$$\|DX_{T^1}(x)/E_x^s\| \leq \frac{K\alpha}{2R} m(DX_t(x)/E_x^c), \quad (6)$$

$\forall x \in \Lambda, \forall t \geq 0$.

Since $E_\Lambda^s \oplus E_\Lambda^c$ is invariant we have $\pi_{X_t(x)}^{s(c)} \circ DX_t(x) = DX_t(x) \circ \pi_x^{s(c)}$, and so

$$\angle(DX_t(x)v_x, E_{X_t(x)}^c) = \frac{\|DX_t(x)v_x^s\|}{\|DX_t(x)v_x^c\|}, \quad \forall x \in \Lambda \quad \forall t \geq 0. \quad (7)$$

Recall that $\angle$ denotes the tangent of the angle. The inequality (6) and the last equality imply

$$\angle(DX_{T^1}(x)v_x, E_{X_{T^1}(x)}^c) \leq \frac{K\alpha}{2R}, \quad \forall x \in \Lambda, \quad \forall v_x \in C_\alpha(E_x^c).$$

So,

$$DX_{T^1}(x)(C_\alpha(E_x^c)) \subset C_{\frac{K\alpha}{2R}}(E_{X_{T^1}(x)}^c), \quad \forall x \in \Lambda.$$

Choose a neighborhood $U^1 = U_\alpha^1 \subset U(\Lambda)$ of Λ sufficiently close to Λ such that

$$DX_{T^1}(x)(C_\alpha(E_x^c)) \subset C_{\frac{K\alpha}{R}}(E_{X_{T^1}(x)}^c), \quad \forall x \in U^1. \quad (8)$$

On the other hand, using (4) we get

$$\frac{\|DX_t(x)v_x^s\|}{\|v_x^s\|} \leq K^{-1}e^{-\lambda t} \frac{\|DX_t(x)v_x^c\|}{\|v_x^c\|}$$

and so,

$$\frac{\|DX_t(x)v_x^s\|}{\|DX_t(x)v_x^c\|} \leq K^{-1}e^{-\lambda t} \frac{\|v_x^s\|}{\|v_x^c\|} = K^{-1}e^{-\lambda t} \angle(v_x, E_x^c).$$

So, by (7), we get

$$\angle(DX_r(x)v_x, E_{X_r(x)}^c) \leq K^{-1}e^{-\lambda r} \angle(v_x, E_x^c) \leq K^{-1} \angle(v_x, E_x^c),$$

$\forall x \in \Lambda, \forall r \in [0, T^1], \forall v_x \in T_x M$. This implies

$$DX_r(x) \left(C_{\frac{K\alpha}{R}}(E_x^c) \right) \subset C_{\frac{\alpha}{R}}(E_{X_r(x)}^c), \quad \forall x \in \Lambda \quad \forall r \in [0, T^1].$$

Choose a neighborhood $V^2 = V_\alpha^2 \subset U^1$ of Λ sufficiently close to Λ such that

$$DX_r(x) \left(C_{\frac{K\alpha}{R}}(E_x^c) \right) \subset C_{\frac{2\alpha}{R}}(E_{X_r(x)}^c), \quad \forall x \in V^2, \quad \forall r \in [0, T^1]. \quad (9)$$

As Λ is an attractor there is a neighborhood $U^2 \subset V^2$ of Λ such that

$$X_t(U^2) \subset V^2, \quad \forall t \geq 0.$$

Now, let $x \in U^2$ and $t \geq T^1$ be given. Then, $t = nT^1 + r$ for some integer $n \geq 1$ and some $r \in [0, T^1]$. Thus,

$$DX_t(x)(C_\alpha(E_x^c)) = DX_{nT^1+r}(x)(C_\alpha(E_x^c)) =$$

$$= DX_r(X_{nT^1}(x))(DX_{nT^1}(x)(C_\alpha(E_x^c))). \quad (10)$$

Using (8) and (10) recursively, and that $\frac{K\alpha}{R} < \alpha$, and $n \geq 1$, we obtain

$$\begin{aligned} DX_{nT^1}(x)(C_\alpha(E_x^c)) &= DX_{(n-1)T^1}(X_{T^1}(x))(DX_{T^1}(x)C_\alpha(E_x^c)) \subset \\ &\subset DX_{(n-1)T^1}(X_{T^1}(x)) \left(C_{\frac{K\alpha}{R}}(E_{X_{T^1}(x)}^c) \right) \subset DX_{(n-1)T^1}(X_{T^1}(x))(C_\alpha(E_{X_{T^1}(x)}^c)) \subset \\ &\subset \cdots \subset DX_{T^1}(X_{(n-1)T^1}(x))(C_\alpha(E_{X_{(n-1)T^1}(x)}^c)) \subset C_{\frac{K\alpha}{R}}(E_{X_{nT^1}(x)}^c). \end{aligned}$$

Henceforth

$$DX_{nT^1}(x)(C_\alpha(E_x^c)) \subset C_{\frac{K\alpha}{R}}(E_{X_{nT^1}(x)}^c).$$

Applying $DX_r(X_{nT^1}(x))$ to both sides of the last expression, replacing in (10) and using (9) we obtain

$$DX_t(x)(C_\alpha(E_x^c)) \subset DX_r(X_{nT^1}(x)) \left(C_{\frac{K\alpha}{R}}(E_{X_{nT^1}(x)}^c) \right) \subset C_{\frac{2\alpha}{R}}(E_{X_t(x)}^c).$$

As $R > 4$ we have $\frac{2\alpha}{R} < \frac{\alpha}{2}$ and so

$$DX_t(x)(C_\alpha(E_x^c)) \subset C_{\frac{\alpha}{2}}(E_{X_t(x)}^c).$$

$\forall x \in U^2, \forall t \geq T^1$, proving (1) of Lemma 3.1.

Throughout we fix the neighborhood U^2 of Λ and the constant $T^1 > 0$ obtained above.

As E_Λ^c is (K, λ) -volume expanding we have

$$\text{vol}(DX_t(x)v_x^c, DX_t(x)w_x^c) \geq Ke^{\lambda t} \text{vol}(v_x^c, w_x^c),$$

$\forall x \in \Lambda, \forall t \geq 0, \forall v_x^c, w_x^c \in E_x^c$.

Clearly there is $L > 1$ such that

$$L^{-1} \cdot \text{vol}(v_x^c, w_x^c) \leq \text{vol}(v_x, w_x) \leq L \cdot \text{vol}(v_x^c, w_x^c),$$

$\forall x \in \Lambda, \forall v_x, w_x \in C_\alpha(E_x^c), \forall \alpha \in (0, 1]$. Applying the last two relations and the invariance of $E_\Lambda^s \oplus E_\Lambda^c$ we obtain

$$\begin{aligned} \text{vol}(DX_t(x)v_x, DX_t(x)w_x) &\geq L^{-1} \text{vol}(DX_t(x)v_x^c, DX_t(x)w_x^c) \geq \\ &\geq L^{-1} Ke^{\lambda t} \text{vol}(v_x^c, w_x^c) \geq L^{-2} Ke^{\lambda t} \text{vol}(v_x, w_x), \end{aligned}$$

$\forall x \in \Lambda, \forall t \geq T^1, \forall v_x, w_x \in C_\alpha(E_x^c)$ (note that $DX_t(x)v_x, DX_t(x)w_x \in C_\alpha(E_{X_t(x)}^c)$ since $t \geq T^1$).

Choose $S > 0$ large so that

$$\frac{S}{L^{-2}K} > 1. \quad (11)$$

It follows that there is $T^2 = T_\alpha^2 > T^1$ such that

$$\text{vol}(DX_{T^2}(x)v_x, DX_{T^2}(x)w_x) \geq \frac{2S}{L^{-2}K} \text{vol}(v_x, w_x),$$

$\forall x \in \Lambda, \forall v_x, w_x \in C_\alpha(E_x^c)$. In particular,

$$\inf\{\text{vol}(DX_{T^2}(x)v_x, DX_{T^2}(x)w_x) : x \in \Lambda,$$

$$v_x, w_x \in C_\alpha(E_x^c), \|v_x\| = \|w_x\| = 1, \langle v_x, w_x \rangle = 0\} \geq \frac{2S}{L^{-2}K}.$$

Since Λ is compact there is a neighborhood $V^3 = V_\alpha^3 \subset U^2$ of Λ so that

$$\inf\{\text{vol}(DX_{T^2}(x)v_x, DX_{T^2}(x)w_x) : x \in U^3,$$

$$v_x, w_x \in C_\alpha(E_x^c), \|v_x\| = \|w_x\| = 1, \langle v_x, w_x \rangle = 0\} \geq \frac{S}{L^{-2}K}.$$

Then,

$$\text{vol}(DX_{T^2}(x)v_x, DX_{T^2}(x)w_x) \geq \frac{S}{L^{-2}K} \text{vol}(v_x, w_x), \quad (12)$$

$\forall x \in U^3, \forall v_x, w_x \in C_\alpha(E_x^c)$ with $\langle v_x, w_x \rangle = 0$. As Λ is an attractor there is a neighborhood $U^3 \subset V^3$ of Λ such that

$$X_t(U^3) \subset V^3, \quad \forall t \geq 0.$$

We have

$$\|P_x^{nT^2}v_x\| \|X(X_{nT^2}(x))\| = \|P_{X_{(n-1)T^2}(x)}^{T^2}(P_x^{(n-1)T^2}v_x\| \|X(X_{T^2}(X_{(n-1)T^2}(x)))\|.$$

Call $z = X_{(n-1)T^2}(x)$, and $v_z = P_x^{(n-1)T^2}v_x$. From the last equality, using that $X(X_{nT^2}(x)) = DX_{nT^2}(x)(X(x))$, v_z is orthogonal to z , and combining (3) with (12) we get

$$\begin{aligned} & \|P_x^{nT^2}v_x\| \|X(X_{nT^2}(x))\| = \|P_z^{T^2}v_z\| \|X(X_{T^2}(z))\| = \\ & \text{vol}(DX_{T^2}(z)v_z, X(X_{T^2}(z))) = \text{vol}(DX_{T^2}(z)v_z, DX_{T^2}(z)(X(z))) \geq \\ & \frac{S}{L^{-2}K} \text{vol}(v_z, X(z)) = \frac{S}{L^{-2}K} \text{vol}(P_x^{(n-1)T^2}v_x, X(X_{(n-1)T^2}(x))). \end{aligned}$$

Thus,

$$\|P_x^{nT^2}v_x\| \cdot \|X(X_{nT^2}(x))\| \geq \left\{ \frac{S}{L^{-2}K} \right\}^n \|v_x\| \cdot \|X(x)\|, \quad (13)$$

$\forall x \in U^3$ regular with $X(x) \in C_\alpha(E_x^c)$, $\forall n \in \mathbb{N}$, $\forall v_x \in C_\alpha(E_x^c) \cap N_x$ (recall that N_x denotes the orthogonal complement of $X(x)$ in $T_x M$).

On the other hand

$$\text{vol}(DX_r(x)v_x, DX_r(x)w_x) \geq L^{-2}K \cdot \text{vol}(v_x, w_x),$$

$$\forall x \in \Lambda, \quad \forall r \in [0, T^2], \quad \forall v_x, w_x \in C_\alpha(E_x^c).$$

As before there is a neighborhood $V^4 = V_\alpha^4 \subset U^3$ of Λ such that

$$\text{vol}(DX_r(x)v_x, DX_r(x)w_x) \geq \frac{L^{-2}K}{2} \text{vol}(v_x, w_x), \quad (14)$$

$\forall x \in V^4$, $\forall v_x, w_x \in C_\alpha(E_x^c)$ with $\langle v_x, w_x \rangle = 0$, $r \in [0, T^2]$. As Λ is an attractor there is a neighborhood $U^4 \subset V^4$ of Λ such that

$$X_t(U_4) \subset V^4, \quad \forall t \geq 0.$$

Now, let $x \in U^4$ regular with $X(x) \in C_\alpha(E_x^c)$, $t \geq T^2$ and $v_x \in C_\alpha(E_x^c) \cap N_x$. Then, $t = nT^2 + r$ for some integer $n \geq 1$ and some $r \in [0, T^2]$.

Applying (13), (14), and using (3) and (11) we obtain

$$\begin{aligned} \|P_x^t v_x\| \cdot \|X(X_t(x))\| &= \|P_{X_{nT^2}(x)}^r P_x^{nT^2} v_x\| \cdot \|DX_r(X_{nT^2}(x))X(X_{nT^2}(x))\| = \\ &= \text{vol}\left(DX_r(X_{nT^2}(x))P_x^{nT^2} v_x, DX_r(X_{nT^2}(x))X(X_{nT^2}(x))\right) \geq \\ &\geq \frac{L^{-2}K}{2} \text{vol}(P_x^{nT^2} v_x, X(X_{nT^2}(x))) = \frac{L^{-2}K}{2} \|P_x^{nT^2} v_x\| \cdot \|X(X_{nT^2}(x))\| \geq \\ &\geq \left(\frac{L^{-2}K}{2}\right) \cdot \left\{ \frac{SN}{L^{-2}K} \right\}^n \cdot \|v_x\| \cdot \|X(x)\| = \\ &= \left(\frac{L^{-2}K}{2}\right) \cdot \left(\frac{S}{L^{-2}K}\right)^{-\frac{r}{T^2}} \left\{ \left(\frac{S}{L^{-2}K}\right)^{\frac{1}{T^2}} \right\}^t \cdot \|v_x\| \cdot \|X(x)\| \geq \\ &\geq \left(\frac{L^{-4}K^2}{2S}\right) \cdot \left\{ \left(\frac{S}{L^{-2}K}\right)^{\frac{1}{T^2}} \right\}^t \cdot \|v_x\| \cdot \|X(x)\|. \end{aligned}$$

Thus, choosing $U_\alpha = U^4$, $T_\alpha = T^2$, $K_\alpha = \frac{L^{-4}K^2}{2S}$ and $\lambda_\alpha = \ln\left(\frac{S}{L^{-2}K}\right)^{\frac{1}{T^2}} > 0$ we obtain (2) of Lemma 3.1. $\square$

Lemma 3.2. *Let Λ a singular-hyperbolic attractor of X , $\alpha > 0$ and $U_\alpha \subset U(\Lambda)$ be a neighborhood of Λ . Let $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$ the splitting of Λ and $E_{U_\alpha}^c$ be a continuous extension of E_Λ^c to U_α . Then, for every singularity σ of X in Λ there are singular cross-sections S^t, S^b associated to σ such that $S^t \cup S^b \subset U_\alpha$,*

$$\Lambda \cap (\partial^h S^t \cup \partial^h S^b) = \emptyset, \quad \text{and} \quad X(x) \in C_\alpha(E_x^c),$$

$$\forall x \in S^t \cup S^b.$$

Proof. Let Λ, α be as in the statement. By Remark 2.1 applied to $U = U_\alpha$ there are singular cross-sections S_0^t, S_0^b associated to σ such that $S_0^t, S_0^b \subset U(\Lambda)$ and $\Lambda \cap (\partial^h S_0^t \cup \partial^h S_0^b) = \emptyset$. Recall that $U(\Lambda)$ is the neighborhood of Λ where the (K, λ) -splitting $T_\Lambda M = E_\Lambda^s \oplus E_\Lambda^c$ has an extension to $T_{U(\Lambda)} M = E_{U(\Lambda)}^s \oplus E_{U(\Lambda)}^c$, with $E_{U(\Lambda)}^s$ invariant and contracting. We denote by l_0^t, l_0^b the singular curves of S_0^t, S_0^b respectively.

Choose two sequences of singular cross-sections $S_n^t \subset S_0^t, S_n^b \subset S_0^b$ associated to σ satisfying

- (a) $\Lambda \cap (\partial^h S_n^t \cup \partial^h S_n^b) = \emptyset$;
- (b) $\text{diam}(S_n^t), \text{diam}(S_n^b) \rightarrow 0$ as $n \rightarrow \infty$;
- (c) the singular curves l_n^t, l_n^b of S_n^t, S_n^b satisfy $l_n^t = l_0^t, l_n^b = l_0^b, \forall n$.

The properties (b) and (c) imply that $\forall n$ large there is $T = T_n > 0$ such that

$$\angle(X(X_T(x)), E_\sigma^c) < \frac{\alpha}{2},$$

$\forall x \in S_n^t \cup S_n^b$. As $E_{U_\alpha}^c$ is a continuous extension of E_Λ^c , we have that $E_{X(X_T(x))}^c$ is close to $E_\sigma^c, \forall x \in S_n^t \cup S_n^b$. Then,

$$X_T(x) \in U_\alpha, \quad \text{and} \quad X(X_T(x)) \in C_\alpha(E_{X(X_T(x))}^c),$$

$\forall n$ large, $\forall x \in S_n^t \cup S_n^b$.

By the Property (a) and the above relation we have that for n large enough $S^t = X_T(S_n^t)$ and $S^b = X_T(S_n^b)$ are singular cross-sections associated to σ satisfying the required properties. $\square$

References

- [1] Bonatti, C.. *C¹-Generic Dynamics: Tame and Wild Behavior*. ICM 2002 (2002) **2**, 265-277.

- [2] Bowen, R.. *Markov partitions and minimal sets for Axiom A diffeomorphisms*. Amer. J. Math. **92** (1970), 907-918.
- [3] Bowen, R.. *Symbolic dynamics for hyperbolic flows*. Amer. J. Math. **95** (1973), 429-460.
- [4] Doering C., I.. Persistently transitive vector fields on three- dimensional manifolds. In *Procs. on Dynamical Systems and Bifurcation Theory*, 1987.
- [5] de Melo, W., Palis, J.. *Geometric Theory of Dynamical Systems-An Introduction*. Springer Verlag, Berlin, 1982.
- [6] Guckenheimer, J., Williams, R., F.. *Structural stability of Lorenz attractors*. Inst. Hautes Études Sci. Publ. Math. **50** (1979), 59-72.
- [7] Hirsch, M., W.. *On invariant subsets of hyperbolic sets*. Essays on Topology and Related Topics, Mémoires dédiés à Georges de Rham (1970), 126-135.
- [8] Hirsch, M., W., Pugh, C., C., Shub, M.. *Invariant manifolds*, volume 583 of *Lect. Notes in Math.* Springer Verlag, Berlin, 1977.
- [9] Hurewicz, W., Wallman, H.. *Dimension Theory*. Princeton University Press, Princeton, NJ, (1984).
- [10] Keynes, H., B., Sears, M.. *Real-expansive flows and topological dimension*. Ergodic Theory Dynamical Systems **1** (1981), 179-195.
- [11] Labarca, R., Pacifico, M., J.. *Stability of singular horseshoes*. Topology **25** (1986), 337-352.
- [12] Mañé, R.. *An ergodic closing lemma*. Ann. of Math. (2) **116** (1982), 503-540.
- [13] Mañé, R.. *Expansive homeomorphisms and topological dimension*. Trans. Amer. Math. Soc. **252** (1979), 313-319.
- [14] Morales, C. A.. Singular-hyperbolic sets and topological dimension. Preprint (2002) to appear.
- [15] Morales, C., A., Pacifico, M., J.. *A dichotomy for three-dimensional vector fields*. To appear in Ergodic Theory Dynamical Systems.
- [16] Morales, C., A., Pacifico, M., J., Pujals, E., R.. *On C^1 robust singular transitive sets for three-dimensional flows*. C. R. Acad. Sci. Paris Sér. I Math. **326** (1998), 81-86.

- [17] Morales, C., A., Pacifico, M., J., Pujals, E., R.. Singular hyperbolic systems. Proc. Amer. Math. Soc., 127, Number 11:3393–3401, 1999.
- [18] Przytycki, F.. *Construction of invariant sets for Anosov diffeomorphisms and hyperbolic attractors*. Studia Math. **68** (1980), 199-213.
- [19] Shilnikov, L., P., Turaev, D., V.. *An example of a wild strange attractor*. (Russian) Mat. Sb. **189** (1998), 137-160; translation in Sb. Math. **189** (1998), 291-314.

C. A. Morales
 Instituto de Matemática
 Universidade Federal do Rio de Janeiro
 C. P. 68.530, CEP 21.945-970
 Rio de Janeiro, R. J. , Brazil
 e-mail: *morales@impa.br*